


# Rx FINASTERIDE

(This medication is used to treat male pattern hair loss)



## USES

This medication is used to treat male pattern hair loss and is to be used only by adult men.

## HOW TO USE

Take one tablet daily or as directed by your doctor. This drug may be taken with or without meals. This drug usually must be taken regularly for 3 months for the benefits to be noticeable. Once treatment is stopped, any gain in hair quantity will be lost. If this medication is crushed prior to administration, it must not be handled by women who are pregnant or who may become pregnant.

## SIDE EFFECTS

Unlikely to occur but the possible side effects include impotence, decreased sex drive or decrease in amount of ejaculate. If you notice other effects not listed above, contact your doctor or pharmacist.

## PRECAUTIONS

Before using this drug, tell your doctor your medical history, especially of: liver disease, prostate cancer, urinary tract disease, drug allergies. The drug can be absorbed through the skin. If the film coating of the tablet has been broken or the tablet crushed, it should not be handled by a woman who is pregnant or plans to become pregnant. Any contact with finasteride by a developing male fetus could result in abnormalities of the external sex organs. Finasteride should not be used in women or children. Finasteride must not be used during pregnancy. If you become pregnant or think you may be pregnant, inform your doctor immediately. It is not known if finasteride is excreted into breast milk. However, this drug is not indicated for use in women.

## DRUG INTERACTIONS

Tell your doctor of all over-the-counter and prescription medication you may use. Do not start or stop any medicine without doctor or pharmacist approval.

## OVERDOSE

If overdose is suspected, contact your local poison control center or emergency room immediately. US residents can call the US national poison hotline at 1-800-222-1222. Canadian residents should call their local poison control center directly.

## MISSED DOSE

If you miss a dose, take it as soon as remembered; do not take it if it is almost time for the next dose, instead, skip the missed dose and resume your usual dosing schedule. Do not "double-up" the dose to catch up.

## STORAGE

Store at room temperature away from sunlight and moisture.

**Note :** This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.



Taj Group of Companies <sup>INDIA</sup>

Taj Pharmaceuticals Ltd.

Phone : General EPA BX : 91 - (0)22 - 26374592/92 91,  
(0)22 - 26374592/93 91 - (0)22 - 30601000,  
Fax : 91-(0)22-26341274

Revised November 2010