



Rx ONDANSETRON

(This medication is used to prevent nausea and vomiting caused by cancer chemotherapy)



USES

This medication is used to prevent nausea and vomiting caused by cancer chemotherapy or after surgery. It works by blocking the hormone (serotonin) that causes vomiting.

OTHER USES

This medication may also be used to treat alcoholism.

HOW TO USE

This medication may be taken with or without food. It may also be taken with antacids. The first dose is usually taken 30 minutes before chemotherapy. Take further doses as directed, usually for 1-2 days after completion of chemotherapy. Other medical conditions and procedures require different dosing schedules. Follow your doctor's orders carefully.

SIDE EFFECTS

Diarrhea or constipation, headache, lightheadedness or drowsiness (uncommon), as well as blurred vision may occur. If these effects persist or worsen, notify your doctor promptly. Very unlikely but report promptly: chest pain. In the unlikely event you have an allergic reaction to this drug, seek immediate medical attention. Symptoms of an allergic reaction include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS

Tell your doctor your medical history, especially of: stomach/intestinal problems (e.g., distension), liver disease, any allergies. Limit alcohol intake, as it may intensify drug side effects. Caution performing tasks requiring alertness (e.g., driving) until you know how this medication affects you. Tell your doctor if you are pregnant before using this medication. It is not known if this drug is excreted into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS

Tell your doctor of all prescription and nonprescription medicines you may use. Do not start or stop any medicine without doctor or pharmacist approval.

OVERDOSE

If overdose is suspected, contact your local poison control center or emergency room immediately. US residents can call the US national poison hotline at 1-800-222-1222. Canadian residents should call their local poison control center directly. Symptoms of overdose may include change in vision, severe constipation, severe dizziness, or irregular heartbeat.

MISSED DOSE

Try to take each dose at the scheduled time. If you miss a dose, take it as soon as remembered; do not take it if it is near the time for the next dose, instead, skip the missed dose and resume your usual dosing schedule. Do not "double-up" the dose to catch up.

STORAGE

Tablets may be stored in or out of the refrigerator between 36 and 86 degrees F (2 and 30 degrees C). Store liquid form between 59 and 86 degrees F (15 and 30 degrees C).

Note : This product information is intended only for residents of the India. Taj Pharmaceuticals Limited, medicines help to treat and prevent a range of conditions—from the most common to the most challenging—for people around the world.



Taj Group of Companies ^{INDIA}

Taj Pharmaceuticals Ltd.

Phone : General EPA BX : 91 - (0)22 - 26374592/92 91,
(0)22 - 26374592/93 91 - (0)22 - 30601000,
Fax : 91-(0)22-26341274

Revised November 2010